



Weekly Local Influenza Surveillance Bulletin


Released December 2, 2009

Region of Waterloo

PUBLIC HEALTH

Influenza Activity: September 1, 2009 to November 28, 2009 (Week 35-47)

Figure 1: Number of laboratory confirmed cases of influenza by week and type, Waterloo Region, September 1, 2009 to November 28, 2009³




No. of Cases by Type	
Influenza A – Total	252
pH1N1	115
H3	0
H3N2	1
Other (non-pH1N1)	0
Not sub-typed ¹	136
Influenza B	0
Total	252

Influenza Activity Level ²	
	No activity
✓	Sporadic
	Localized
	Widespread

Updated: 2-Dec-09

Figure 2: Number of laboratory confirmed cases of influenza by age and type, Waterloo Region, September 1, 2009 to November 28, 2009




¹ Not sub-typed includes all Influenza A cases that had a sub-type of "blank", "untypeable" or "not-subtyped".

² For definitions of influenza activity levels: <http://www.phac-aspc.gc.ca/fluwatch/09-10/def09-10-eng.php>

³ Note: In Ontario, the onset date of symptoms for the first confirmed case of Influenza A pH1N1 virus was April 11, 2009. The weeks refer to FluWatch weeks, which are established by the Public Health Agency of Canada.

Figure 3: Laboratory confirmed cases of pandemic H1N1 in Ontario by age group and wave, April 13, 2009 to November 21, 2009



SOURCE: Ontario Ministry of Health and Long-Term Care, integrated Public Health Information System (iPHIS) database, extracted [25/11/2009].

Figure 4: Number of laboratory confirmed cases of influenza A by week, Waterloo Region, 2005-2009




Table 1: Pandemic H1N1 indicators, Waterloo Region and Ontario, Sept. 1, 2009 - Nov. 28, 2009

Indicator	Waterloo Region*	Ontario*
Laboratory confirmed H1N1 cases	115	4183
Number of H1N1 hospitalizations	22	1158
Population-based hospitalization rate	4.4 per 100,000 residents	8.9 per 100,000 Ontarians
Number of deaths	3	70
Population-based mortality rate	0.60 per 100,000 residents	0.54 per 100,000 Ontarians
Age of laboratory confirmed H1N1 cases [^]	<1 – 74 years	<1 – 90+ years
Age of hospitalized cases	Range: <1 – 74 years Median age: 14 years Average age: 26 years	Range: 0 – 90 years Median age: 24 years Average age: 29 years
Age of fatal cases	Range: >35 years	Range: 0 – 95 years Median age: 54 years Average age: 51 years

* Waterloo Region data is current as of November 28, 2009, while Ontario data is current as of November 21, 2009, except for hospitalizations and deaths, which are current as of November 25, 2009


[^] Ontario data reflects all cases from Week 14-46 (April 5, 2009 to November 21, 2009).

Proxy Indicators for Influenza Surveillance

Influenza-Like-Illness

The number of Influenza-Like Illness (ILI)⁵ visits is a useful indicator of the level of community transmission of pH1N1 and the resulting burden on primary care practices. Transmission patterns suggest that pH1N1 is more prevalent in the younger age groups (5 – 19 years). As such, we are tracking the ILI consultation rate at post-secondary institutions⁶ health services in Waterloo Region as well as in select primary care practices⁷. ILI information may change on a weekly basis as data is collected from additional primary care practices. Therefore, rates reported in the current bulletin may not equal rates reported in previous weeks.

Figure 5: Influenza-Like-Illness (ILI) consultations per 1,000 patient visits, Waterloo and Ontario, September 13 – November 28, 2009



⁵ Case definition for ILI is: acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which could be due to influenza virus. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

⁶ Post-secondary institutions include Conestoga College, Wilfrid Laurier University, and University of Waterloo

⁷ Data from an additional primary care practice was included during week 43.

Hospital-Based Data

The Ministry of Health and Long Term Care (MOHLTC) is tracking the number of Emergency Room visits and the number of hospital-based deaths as useful indicators of the level of community transmission of pH1N1 and the resulting burden on hospitals. The figures below include aggregated data from all three acute care hospitals within the Region of Waterloo: Cambridge Memorial Hospital, Grand River Hospital, and St. Mary's General Hospital.

Figure 6: Total all-cause Emergency Room (ER) visits by week, Waterloo Region, September 6 – November 28, 2009






Figure 7: Total all-cause hospital-based mortality by week, Waterloo Region, September 6 – November 28, 2009



School Absenteeism

Since pH1N1 is more prevalent in the younger age groups, monitoring school absenteeism rates is a useful indicator of the level of community transmission of the virus. An absenteeism rate of 10% or greater has been identified as a threshold by the MOHLTC to signify higher than expected levels of school absenteeism. The figure below shows the average number and average proportion of schools in the Region of Waterloo that meet this threshold, by week and type of school (elementary or secondary).

Figure 8: Average number and average proportion of schools reporting $\geq 10\%$ absenteeism (all-cause) by week and type of school, Waterloo Region, October 6 – November 28, 2009



For provincial and national influenza information, please visit the following websites:

MINISTRY OF HEALTH AND LONG-TERM CARE (MOHLTC)

The latest Ontario Influenza Bulletin can be viewed at the following site:

http://www.health.gov.on.ca/english/providers/program/pubhealth/flu/flu_08/flubul_mn.html

The latest information on Influenza A pH1N1 virus in Ontario can be viewed at the following site:

http://www.health.gov.on.ca/english/public/updates/archives/hu_09/provider/default.html

HEALTH CANADA

The latest Health Canada FluWatch can be viewed at the following site:

<http://www.phac-aspc.gc.ca/fluwatch/08-09/index-eng.php>